

Introducing the NEW Standard for Early Blight Control in Potatoes

Provysol™ Fungicide Guidelines for Use in Potatoes

- **Active Ingredient:** Mefentrifluconazole (FRAC 3)
- **Crop:** Potatoes
- **Target Disease:** Early Blight, Brown Spot, Black Dot
- **Optimal Timing:** Crop row closure, prior to onset of disease
- **Use Rate:** 3-5 oz/A
- **Application Method:** Ground, Air, or Chemigation
- **Optimal Spray Volume:** Enough for complete coverage. 20 GPA (ground) or 5 GPA (aerial).
- **Tank-Mix:** Can be tank-mixed with most commonly-used white mold and late blight fungicides
- **Re-Entry Interval:** 12 hours
- **Preharvest Interval:** 7 days

This quick guide DOES NOT replace the need for a product label.
It is designed to supplement the product label.

Always read and follow label directions.

Provysol is a trademark of BASF. ©2019 BASF Corporation. All rights reserved.



BASF
We create chemistry

Priaxor®

Xemium® Brand Fungicide

Serifel®

Biofungicide

Outlook®

Herbicide



Zidua® SC

Herbicide

Regent® 4 SC

Insecticide

Priaxor®

Xemium® Brand Fungicide

Fastac® CS

Insecticide

OR

Sefina®

Inscalis® Insecticide

Rely²⁸⁰

Herbicide

Provysol™

Fungicide

New for 2020!

Endura®

Fungicide

Serifel®

Biofungicide

Forum®

Fungicide

Prowl® H2O

Herbicide

Planting

Sprout Development

Vegetative

Bulking

Maturation

Harvest

